

author: Imre Rábai

In honor of the Janus Pannonius High School in Pécs, where I taught in the 1956–57 school year.

translated from: KoMaL, February 1999, page 87

Pdf file: KoMaL, February 1999, page 87

In 1983, a new system of entrance examination was adopted. In order to be accepted to a university, the 3rd and 4th years of high school grades in mathematics, Hungarian language and literature, history, foreign language, physics, (biology, chemistry, geography, another foreign language - students choose from these) are counted toward university entrance performance.

The 'brought' points (i.e. the points comprised of final grades in high school in the subjects listed above) add up to a total of 60 points. In addition, students are assessed by written and oral examinations for a total of 60 points. So, there is a total of 120 points possible.

In Mathematics, the same exam serves as the GED and the university entrance exam. These problem sets consist of 8 problems, presented in order of difficulty (from easiest to most difficult).

This problem set is similar to such an exam. We advise the reader to work through the problem set while measuring the time. There are 180 minutes to solve and present all problems.

1. Solve the following equations over the real numbers.

$$\text{a) } 2^x + \sqrt{2^{x+2} \cdot 5^x} = 3 \cdot 5^x \quad \text{b) } \frac{1}{x^2 - 3x + 3} + \frac{2}{x^2 - 3x + 4} = \frac{6}{x^2 - 3x + 5} \quad \text{c) } 2 \sin 2x + 2 \sin x = 2 \cos x + 1$$

Solution: a) $\sqrt{2^{x+2} 5^x} = 2 \cdot 2^{\frac{x}{2}} 5^{\frac{x}{2}}$. Dividing by 5^x and writing $u = \left(\frac{2}{5}\right)^{\frac{x}{2}} > 0$: $u^2 + 2u - 3 = 0$, so $u = 1$ (the root -3 is negative), and $x = 0$.

b) Let $y = x^2 - 3x + 4 = \left(x - \frac{3}{2}\right)^2 + \frac{7}{4} \geq \frac{7}{4}$. The equation is $\frac{1}{y-1} + \frac{2}{y} = \frac{6}{y+1}$, which gives $3y^2 - 7y + 2 = 0$, so $y = 2$ or $y = \frac{1}{3}$; only $y = 2$ is possible. Then $x^2 - 3x + 2 = 0$, and the solutions are $x = 1$ and $x = 2$.

c) $4 \sin x \cos x + 2 \sin x - 2 \cos x - 1 = 2 \sin x (2 \cos x + 1) - (2 \cos x + 1) = (2 \cos x + 1) (2 \sin x - 1) = 0$. So $\cos x = -\frac{1}{2}$ or $\sin x = \frac{1}{2}$:

$$x = \pm \frac{2\pi}{3} + 2k\pi, \quad x = \frac{\pi}{6} + 2k\pi, \quad x = \frac{5\pi}{6} + 2k\pi, \quad k \text{ is an integer}$$

2. Solve the following inequalities over the real numbers.

$$\text{a) } (x+1)(|x|-1) \leq -\frac{1}{2} \quad \text{b) } \tan x + \frac{1}{\tan x} \geq -2 \quad \text{c) } \log_2(4^x - 5 \cdot 2^x + 8) > 2$$

Solution: a) If $x \geq 0$: $x^2 - 1 \leq -\frac{1}{2}$, so $0 \leq x \leq \frac{1}{\sqrt{2}}$. If $x < 0$: $-(x+1)^2 \leq -\frac{1}{2}$, that is, $|x+1| \geq \frac{1}{\sqrt{2}}$. The solution is

$$x \leq -1 - \frac{\sqrt{2}}{2} \quad \text{or} \quad -1 + \frac{\sqrt{2}}{2} \leq x \leq \frac{\sqrt{2}}{2}$$

b) With $t = \tan x \neq 0$: $t + \frac{1}{t} + 2 = \frac{(t+1)^2}{t} \geq 0$, so $t > 0$ or $t = -1$: $k\pi < x < \frac{\pi}{2} + k\pi$, or $x = -\frac{\pi}{4} + k\pi$.

c) $4^x - 5 \cdot 2^x + 8 > 4$, that is, $(2^x - 1)(2^x - 4) > 0$, so $2^x < 1$ or $2^x > 4$: $x < 0$ or $x > 2$.

3. Consider the line e with equation $4x - 3y + 20 = 0$ and its point E with first coordinate $x_0 = -2$. The circle k has radius 10 and is tangent to the line e at the point E .

- a) Find the coordinates of the vectors $\overrightarrow{EK}$ and $\overrightarrow{OK}$, where K is the center of the circle k and O is the origin.
b) Write down the equation of the circle k .

Solution: $E = (-2, 4)$. The vector $\overrightarrow{EK}$ is perpendicular to e and has length 10; the normal vector $(4, -3)$ has length 5. So there are two circles:

a) $\overrightarrow{EK}_1 = (8, -6)$, $\overrightarrow{OK}_1 = (6, -2)$; or $\overrightarrow{EK}_2 = (-8, 6)$, $\overrightarrow{OK}_2 = (-10, 10)$.

b) $(x - 6)^2 + (y + 2)^2 = 100$ or $(x + 10)^2 + (y - 10)^2 = 100$.

4. Prove that the ratio of the two roots of the equation $ax^2 + bx + c = 0$ ($a \neq 0$, $b^2 - 4ac > 0$) is 3 : 2 if and only if $6b^2 = 25ac$.

Solution: If the roots are $3t$ and $2t$, then by Viète's formulas $5t = -\frac{b}{a}$ and $6t^2 = \frac{c}{a}$, so $6 \cdot \frac{b^2}{25a^2} = \frac{c}{a}$, that is, $6b^2 = 25ac$.

Conversely, if $6b^2 = 25ac$, then $b^2 - 4ac = b^2 - \frac{24}{25}b^2 = \left(\frac{b}{5}\right)^2$, and the roots are $\frac{-b \pm \frac{b}{5}}{2a}$, that is, $-\frac{2b}{5a}$ and $-\frac{3b}{5a}$, whose ratio is 3 : 2 (note $b \neq 0$, since the discriminant is positive).

5. In the rectangle $ABCD$, $2 \cdot AB = 3 \cdot BC$. The distances of the vertices C and D from a point P of the line AB are $8\sqrt{2}$ and $4\sqrt{5}$ units. Find the sides of the rectangle. Where is the point P ?

Solution: Let $AB = 3a$, $BC = 2a$, and $A(0, 0)$, $B(3a, 0)$, $C(3a, 2a)$, $D(0, 2a)$, $P(p, 0)$. Since $PC < PD$, P is closer to B :

$$(3a - p)^2 + 4a^2 = 80, \quad p^2 + 4a^2 = 128$$

Subtracting: $9a^2 - 6ap = -48$, so $p = \frac{3a^2 + 16}{2a}$; substituting, $25a^4 - 416a^2 + 256 = 0$, so $a^2 = 16$ or $a^2 = \frac{16}{25}$.

If $a = 4$: the sides are $AB = 12$ and $BC = 8$, and $p = 8$: P is on the side AB , 8 units from A and 4 units from B .

If $a = \frac{4}{5}$: the sides are $AB = \frac{12}{5}$ and $BC = \frac{8}{5}$, and $p = \frac{56}{5}$: P is on the extension of AB beyond B , with $BP = \frac{44}{5}$.

6. Prove that a triangle is a right triangle if and only if $\sin \gamma = \cos \alpha + \cos \beta$, where γ is an acute angle of the triangle.

Solution: If the triangle has a right angle, it is at A or B (since γ is acute); say $\alpha = 90^\circ$. Then $\cos \alpha = 0$ and $\beta = 90^\circ - \gamma$, so $\cos \beta = \sin \gamma$, and the equation holds. The case $\beta = 90^\circ$ is the same.

Conversely, by the sum-to-product formula, and since $\frac{\alpha + \beta}{2} = 90^\circ - \frac{\gamma}{2}$,

$$\cos \alpha + \cos \beta = 2 \cos \frac{\alpha + \beta}{2} \cos \frac{\alpha - \beta}{2} = 2 \sin \frac{\gamma}{2} \cos \frac{\alpha - \beta}{2}$$

and $\sin \gamma = 2 \sin \frac{\gamma}{2} \cos \frac{\gamma}{2}$. Dividing by $2 \sin \frac{\gamma}{2} > 0$: $\cos \frac{\gamma}{2} = \cos \frac{\alpha - \beta}{2}$. Both angles are between -90° and 90° , so $\gamma = \pm(\alpha - \beta)$. If $\gamma = \alpha - \beta$, then $\alpha = \beta + \gamma = 180^\circ - \alpha$, so $\alpha = 90^\circ$; similarly, if $\gamma = \beta - \alpha$, then $\beta = 90^\circ$.

7. Solve the inequality $x^{1+\log_a x} > a^2 x$, where $a > 0$, $a \neq 1$ is a real parameter.

Solution: We need $x > 0$. Let $L = \log_a x$; then $\log_a (x^{1+L}) = (1+L)L$ and $\log_a (a^2 x) = 2 + L$.

If $a > 1$: $\log_a$ is increasing, so $(1+L)L > 2 + L$, that is, $L^2 > 2$: $x > a^{\sqrt{2}}$ or $0 < x < a^{-\sqrt{2}}$.

If $0 < a < 1$: $\log_a$ is decreasing, so $L^2 < 2$: $a^{\sqrt{2}} < x < a^{-\sqrt{2}}$.

8. Find the smallest value of the expression

$$f(x, y) = x^4 + 4y^4 + \frac{1}{x^2y^2}$$

For which pairs (x, y) does $f(x, y)$ take its smallest value?

Solution: We need $xy \neq 0$. By the inequality between the arithmetic and geometric means (twice),

$$x^4 + 4y^4 + \frac{1}{x^2y^2} \geq 2\sqrt{4x^4y^4} + \frac{1}{x^2y^2} = 4x^2y^2 + \frac{1}{x^2y^2} \geq 2\sqrt{4} = 4$$

Equality holds if $x^4 = 4y^4$ and $4x^2y^2 = \frac{1}{x^2y^2}$, that is, $x^2 = 2y^2$ and $x^2y^2 = \frac{1}{2}$: $x^2 = 1, y^2 = \frac{1}{2}$. The smallest value is 4, at the four pairs $\left(\pm 1, \pm \frac{\sqrt{2}}{2}\right)$.