

One important achievement of our studies is the ability to solve quadratic equations with any real coefficients. In what follows, we will master this important skill. As a matter of fact, we can apply the method of completing the square to any quadratic expression. The steps are still the same - it's just that each step takes a bit more work.

Recall that a **perfect square** is the square of an integer. For example, 4, 25, or 49 are perfect squares.

Recall that if N is a non-negative number, then \sqrt{N} is the non-negative number x such that $x^2 = N$. For example, $\sqrt{7}$ is the non-negative number, that, when we square, we get 7. In short, $(\sqrt{7})^2 = 7$. In what follows, we start interpreting numbers that are not perfect squares as squares anyway. For example, we can interpret 7 as $(\sqrt{7})^2$. This will be useful when we want to apply the difference of squares theorem. The expression $x^2 - 4$ is factorable over the integers since $x^2 - 4 = (x + 2)(x - 2)$. On the other hand, $x^2 - 7$ is not factorable over the integers. Over the real numbers, $x^2 - 7$ can be factored as $x^2 - 7 = x^2 - (\sqrt{7})^2 = (x + \sqrt{7})(x - \sqrt{7})$.

Example 1. Solve the equation $x^2 + 13 = 8x$ over the real numbers. Make sure to check your solutions.

Solution: Since the equation is quadratic, we reduce one side to zero and factor the other side by completing the square.

$$\begin{aligned} x^2 + 13 &= 8x \\ x^2 - 8x + 13 &= 0 && (x - 4)^2 = x^2 - 8x + 16 \quad \text{so we smuggle in 16} \\ \underbrace{x^2 - 8x + 16} - 16 + 13 &= 0 \\ (x - 4)^2 - 3 &= 0 && 3 = (\sqrt{3})^2 \\ (x - 4)^2 - (\sqrt{3})^2 &= 0 && \text{factor via difference of squares theorem} \\ (x - 4 + \sqrt{3})(x - 4 - \sqrt{3}) &= 0 \end{aligned}$$

We apply the zero product rule, and solve for the zeroes of each linear factor.

$$\begin{aligned} x - 4 + \sqrt{3} &= 0 && \text{or} && x - 4 - \sqrt{3} &= 0 \\ x + \sqrt{3} &= 4 && && x - \sqrt{3} &= 4 \\ x_1 &= 4 - \sqrt{3} && && x_2 &= 4 + \sqrt{3} \end{aligned}$$

$$x_1 = 4 - \sqrt{3} \quad x_2 = 4 + \sqrt{3} \quad \text{or in shorter notation: } x_{1,2} = 4 \pm \sqrt{3}$$

We check: if $x = 4 - \sqrt{3}$, then

$$\begin{aligned} \text{LHS} &= (4 - \sqrt{3})^2 + 13 = 16 - 4\sqrt{3} - 4\sqrt{3} + 3 + 13 = 32 - 8\sqrt{3} \\ \text{RHS} &= 8(4 - \sqrt{3}) = 32 - 8\sqrt{3} \end{aligned}$$

If $x = 4 + \sqrt{3}$, then

$$\begin{aligned} \text{LHS} &= (4 + \sqrt{3})^2 + 13 = 16 + 4\sqrt{3} + 4\sqrt{3} + 3 + 13 = 32 + 8\sqrt{3} \\ \text{RHS} &= 8(4 + \sqrt{3}) = 32 + 8\sqrt{3} \end{aligned}$$

Thus our solution, $\boxed{4 + \sqrt{3} \text{ and } 4 - \sqrt{3}}$ is correct.

Our solutions are irrational numbers. When a quadratic equation has irrational solutions, all other methods break down, and only completing the square works. If the reader is familiar with methods such as grouping or the AC-method, or trial and

error, try solving $x^2 + 13 = 8x$ using any of those methods!

Being able to handle irrational solutions is a great advantage of the method of completing the square.

Please also note that in case of irrational solutions, writing the exact value necessitates the use of radicals. Recall that the exact value of an irrational number can not be captured as a fraction or as a terminating or repeating decimal, so radical form is our only option.

Example 2. Solve the equation $x^2 = 4x + 1$ over the real numbers. Make sure to check your solutions.

Solution: We complete the square.

$$\begin{aligned}
 x^2 &= 4x + 1 \\
 x^2 - 4x - 1 &= 0 & (x-2)^2 &= x^2 - 4x + 4 \\
 \underbrace{x^2 - 4x + 4}_{(x-2)^2} - 4 - 1 &= 0 \\
 (x-2)^2 - 5 &= 0 & 5 &= (\sqrt{5})^2 \\
 (x-2)^2 - (\sqrt{5})^2 &= 0 \\
 (x-2+\sqrt{5})(x-2-\sqrt{5}) &= 0
 \end{aligned}$$

We apply the zero product rule to the product of the two linear factors.

$$\begin{aligned}
 x - 2 + \sqrt{5} &= 0 & \text{or} & & x - 2 - \sqrt{5} &= 0 \\
 x + \sqrt{5} &= 2 & & & x - \sqrt{5} &= 2 \\
 x_1 &= 2 - \sqrt{5} & & & x_2 &= 2 + \sqrt{5} \\
 x_1 = 2 - \sqrt{5} & \text{ and } & x_2 = 2 + \sqrt{5} & \text{ or in shorter notation: } & x_{1,2} &= 2 \pm \sqrt{5}
 \end{aligned}$$

We check: if $x = 2 - \sqrt{5}$, then

$$\begin{aligned}
 \text{LHS} &= (2 - \sqrt{5})^2 = 4 - 2\sqrt{5} - 2\sqrt{5} + 5 = 9 - 4\sqrt{5} \\
 \text{RHS} &= 4(2 - \sqrt{5}) + 1 = 8 - 4\sqrt{5} + 1 = 9 - 4\sqrt{5}
 \end{aligned}$$

If $x = 2 + \sqrt{5}$, then

$$\begin{aligned}
 \text{LHS} &= (2 + \sqrt{5})^2 = 4 + 2\sqrt{5} + 2\sqrt{5} + 5 = 9 + 4\sqrt{5} \\
 \text{RHS} &= 4(2 + \sqrt{5}) + 1 = 8 + 4\sqrt{5} + 1 = 9 + 4\sqrt{5}
 \end{aligned}$$

Thus our solution, $\boxed{2 + \sqrt{5} \text{ and } 2 - \sqrt{5}}$ is correct.

Example 3. Solve the equation $2x^2 + 10x + 9 = 0$ over the real numbers. Make sure to check your solutions.

Solution: We will factor by completing the square. We first factor out the leading coefficient.

$$\begin{aligned} 2x^2 + 10x + 9 &= 0 \\ 2\left(x^2 + 5x + \frac{9}{2}\right) &= 0 \end{aligned}$$

Half of the linear coefficient is $\frac{5}{2}$. Thus the complete square is $\left(x + \frac{5}{2}\right)^2$

$$\left(x + \frac{5}{2}\right)^2 = \left(x + \frac{5}{2}\right)\left(x + \frac{5}{2}\right) = x^2 + \frac{5}{2}x + \frac{5}{2}x + \frac{25}{4} = x^2 + 5x + \frac{25}{4}$$

So we smuggle in $\frac{25}{4}$.

$$\begin{aligned} 2\left(\underbrace{x^2 + 5x + \frac{25}{4}} - \frac{25}{4} + \frac{9}{2}\right) &= 0 & \frac{9}{2} &= \frac{18}{4} \\ 2\left(\left(x + \frac{5}{2}\right)^2 - \frac{25}{4} + \frac{18}{4}\right) &= 0 \\ 2\left(\left(x + \frac{5}{2}\right)^2 - \frac{7}{4}\right) &= 0 \end{aligned}$$

We next apply the difference of squares theorem. $\frac{7}{4}$ is the square of a number - its own square root.

$$\begin{aligned} 2\left(\left(x + \frac{5}{2}\right)^2 - \frac{7}{4}\right) &= 0 & \frac{7}{4} &= \left(\sqrt{\frac{7}{4}}\right)^2 = \left(\frac{\sqrt{7}}{\sqrt{4}}\right)^2 = \left(\frac{\sqrt{7}}{2}\right)^2 \\ 2\left(\left(x + \frac{5}{2}\right)^2 - \left(\frac{\sqrt{7}}{2}\right)^2\right) &= 0 \\ 2\left(x + \frac{5}{2} + \frac{\sqrt{7}}{2}\right)\left(x + \frac{5}{2} - \frac{\sqrt{7}}{2}\right) &= 0 \end{aligned}$$

We solve for the zeroes of both linear factors:

$$\begin{aligned} x + \frac{5}{2} + \frac{\sqrt{7}}{2} &= 0 & \text{or} & & x + \frac{5}{2} - \frac{\sqrt{7}}{2} &= 0 \\ x + \frac{\sqrt{7}}{2} &= -\frac{5}{2} & & & x - \frac{\sqrt{7}}{2} &= -\frac{5}{2} \\ x_1 &= -\frac{5}{2} - \frac{\sqrt{7}}{2} = \frac{-5 - \sqrt{7}}{2} & & & x_2 &= -\frac{5}{2} + \frac{\sqrt{7}}{2} = \frac{-5 + \sqrt{7}}{2} \end{aligned}$$

So the answer is $x_1 = \frac{-5 - \sqrt{7}}{2}$ and $x_2 = \frac{-5 + \sqrt{7}}{2}$ or, in a more compact form, $x_{1,2} = \frac{-5 \pm \sqrt{7}}{2}$.

We check: if $x = \frac{-5 - \sqrt{7}}{2}$, then

$$\begin{aligned} \text{LHS} &= 2x^2 + 10x + 9 = 2 \left(\frac{-5 - \sqrt{7}}{2} \right)^2 + 10 \left(\frac{-5 - \sqrt{7}}{2} \right) + 9 = \frac{2}{1} \cdot \frac{(-5 - \sqrt{7})^2}{2^2} + \frac{10}{1} \cdot \frac{-5 - \sqrt{7}}{2} + 9 \\ &= \frac{2(-5 - \sqrt{7})^2}{4} + \frac{10(-5 - \sqrt{7})}{2} + \frac{9}{1} = \frac{(-5 - \sqrt{7})^2}{2} + \frac{10(-5 - \sqrt{7})}{2} + \frac{18}{2} = \\ &= \frac{25 + 7 + 10\sqrt{7} - 50 - 10\sqrt{7} + 18}{2} = \frac{32 + 10\sqrt{7} - 50 - 10\sqrt{7} + 18}{2} = 0 = \text{RHS} \end{aligned}$$

and if $x = \frac{-5 + \sqrt{7}}{2}$, then

$$\begin{aligned} \text{LHS} &= 2x^2 + 10x + 9 = 2 \left(\frac{-5 + \sqrt{7}}{2} \right)^2 + 10 \left(\frac{-5 + \sqrt{7}}{2} \right) + 9 = \frac{2}{1} \cdot \frac{(-5 + \sqrt{7})^2}{2^2} + \frac{10}{1} \cdot \frac{-5 + \sqrt{7}}{2} + 9 \\ &= \frac{2(-5 + \sqrt{7})^2}{4} + \frac{10(-5 + \sqrt{7})}{2} + \frac{9}{1} = \frac{(-5 + \sqrt{7})^2}{2} + \frac{10(-5 + \sqrt{7})}{2} + \frac{18}{2} = \\ &= \frac{25 + 7 - 10\sqrt{7} - 50 + 10\sqrt{7} + 18}{2} = \frac{32 - 10\sqrt{7} - 50 + 10\sqrt{7} + 18}{2} = 0 = \text{RHS} \end{aligned}$$

Thus our solution, $\frac{-5 - \sqrt{7}}{2}$ and $\frac{-5 + \sqrt{7}}{2}$ is correct.

Example 4. Is there a real number that is exactly one less than its own square?

Solution: Let us denote this number by x . The square of x is then x^2 . We set up and solve the equation comparing the number to its own square.

$$x = x^2 - 1$$

Since the equation is quadratic, we reduce one side to zero and factor the other side.

$$\begin{aligned} x &= x^2 - 1 \\ 0 &= x^2 - x - 1 \\ x^2 - x - 1 &= 0 & \left(x - \frac{1}{2}\right)^2 &= x^2 - x + \frac{1}{4} \\ \underbrace{x^2 - x + \frac{1}{4}} - \frac{1}{4} - 1 &= 0 \\ \left(x - \frac{1}{2}\right)^2 - \frac{1}{4} - \frac{4}{4} &= 0 \\ \left(x - \frac{1}{2}\right)^2 - \frac{5}{4} &= 0 & \frac{5}{4} &= \left(\sqrt{\frac{5}{4}}\right)^2 = \left(\frac{\sqrt{5}}{\sqrt{4}}\right)^2 = \left(\frac{\sqrt{5}}{2}\right)^2 \\ \left(x - \frac{1}{2}\right)^2 - \left(\frac{\sqrt{5}}{2}\right)^2 &= 0 \\ \left(x - \frac{1}{2} + \frac{\sqrt{5}}{2}\right) \left(x - \frac{1}{2} - \frac{\sqrt{5}}{2}\right) &= 0 \end{aligned}$$

$$\left(x - \frac{1}{2} + \frac{\sqrt{5}}{2}\right) \left(x - \frac{1}{2} - \frac{\sqrt{5}}{2}\right) = 0$$

$$\begin{aligned} \text{Either } x - \frac{1}{2} + \frac{\sqrt{5}}{2} &= 0 & \text{or } x - \frac{1}{2} - \frac{\sqrt{5}}{2} &= 0 \\ x + \frac{\sqrt{5}}{2} &= \frac{1}{2} & x - \frac{\sqrt{5}}{2} &= \frac{1}{2} \\ x_1 &= \frac{1}{2} - \frac{\sqrt{5}}{2} & x_2 &= \frac{1}{2} + \frac{\sqrt{5}}{2} \end{aligned}$$

The greater number, $x_2 = \frac{1 + \sqrt{5}}{2} \approx 1.618034$ is a famous number. It is called the **golden mean** or **golden ratio**. $\frac{1 + \sqrt{5}}{2}$ is often denoted by φ and has very interesting properties. For example, it is one less than its own square.

$$\begin{aligned} \varphi^2 - 1 &= \left(\frac{1 + \sqrt{5}}{2}\right)^2 - 1 = \frac{(\sqrt{5} + 1)^2}{2^2} - 1 = \frac{5 + 1 + 2\sqrt{5}}{4} - 1 = \frac{6 + 2\sqrt{5}}{4} - 1 = \frac{2(3 + \sqrt{5})}{2 \cdot 2} - 1 \\ &= \frac{3 + \sqrt{5}}{2} - \frac{2}{2} = \frac{3 + \sqrt{5}}{2} = \varphi \quad \checkmark \quad \text{and so } \varphi = \frac{1 + \sqrt{5}}{2} \text{ works.} \end{aligned}$$

Let us check the other number. Let us denote $\frac{1 - \sqrt{5}}{2}$ by ψ .

$$\begin{aligned} \psi^2 - 1 &= \left(\frac{1 - \sqrt{5}}{2}\right)^2 - 1 = \frac{(\sqrt{5} - 1)^2}{2^2} - 1 = \frac{5 + 1 - 2\sqrt{5}}{4} - 1 = \frac{6 - 2\sqrt{5}}{4} - 1 = \frac{2(3 - \sqrt{5})}{2 \cdot 2} - 1 \\ &= \frac{3 - \sqrt{5}}{2} - \frac{2}{2} = \frac{3 - \sqrt{5}}{2} = \psi \quad \checkmark \quad \text{and so } \psi = \frac{1 - \sqrt{5}}{2} \text{ also works.} \end{aligned}$$



Sample Problems

1. Solve the equation $3x^2 - 5x - 7 = 0$ over the real numbers. Make sure to check your solutions.
2. Find all numbers that are exactly three less than their own square.



Practice Problems

Solve each of the following equations over the real numbers. Make sure to check your solutions.

- | | | |
|----------------------|--------------------------|------------------------|
| 1. $x^2 - 13 = 6x$ | 4. $3x^2 - 11x + 5 = 0$ | 7. $x^2 + 8x - 10 = 0$ |
| 2. $x^2 = 10x + 3$ | 5. $-2x^2 + 8x - 29 = 0$ | 8. $3x^2 + x - 1 = 0$ |
| 3. $x^2 + x - 1 = 0$ | 6. $5x^2 - 4x = 20$ | 9. $x^2 + x + 1 = 0$ |

10. Find all numbers with the following property: twice the number is two less than the square of the number.
11. Find all numbers with the following property: the number is four less than the square of the number.



Answers

Sample Problems

$$1. \frac{5 - \sqrt{109}}{6} \text{ and } \frac{5 + \sqrt{109}}{6} \quad 2. \frac{1 + \sqrt{13}}{2} \text{ and } \frac{1 - \sqrt{13}}{2}$$

Practice Problems

$$1. 3 \pm \sqrt{22} \quad 2. 5 \pm 2\sqrt{7} \quad 3. \frac{-1 \pm \sqrt{5}}{2} \quad 4. \frac{11 \pm \sqrt{61}}{6} \quad 5. \text{no real solution} \quad 6. \frac{2 \pm 2\sqrt{26}}{5}$$

$$7. -4 \pm \sqrt{26} \quad 8. \frac{-1 \pm \sqrt{13}}{6} \quad 9. \text{no real solution} \quad 10. 1 + \sqrt{3} \text{ and } 1 - \sqrt{3} \quad 11. \frac{1 + \sqrt{17}}{2} \text{ and } \frac{1 - \sqrt{17}}{2}$$

Sample Problems



Solutions

1. Solve the equation $3x^2 - 5x - 7 = 0$ over the real numbers. Make sure to check your solutions.

Solution: We factor out the leading coefficient and factor by completing the square.

$$3x^2 - 5x - 7 = 0$$

$$3 \left(x^2 - \frac{5}{3}x - \frac{7}{3} \right) = 0$$

Half of the linear coefficient is $\frac{5}{3} \div 2 = \frac{5}{3} \cdot \frac{1}{2} = \frac{5}{6}$. Thus the complete square is $\left(x - \frac{5}{6}\right)^2$

$$\left(x - \frac{5}{6}\right)^2 = \left(x - \frac{5}{6}\right) \left(x - \frac{5}{6}\right) = x^2 - \frac{5}{6}x - \frac{5}{6}x + \frac{25}{36} = x^2 - \frac{5}{3}x + \frac{25}{36} \quad \text{So we smuggle in } \frac{25}{36}$$

$$3 \left(\underbrace{x^2 - \frac{5}{3}x + \frac{25}{36}}_{\left(x - \frac{5}{6}\right)^2} - \frac{25}{36} - \frac{7}{3} \right) = 0$$

$$3 \left(\left(x - \frac{5}{6}\right)^2 - \frac{25}{36} - \frac{7 \cdot 12}{3 \cdot 12} \right) = 0$$

$$3 \left(\left(x - \frac{5}{6}\right)^2 - \frac{25}{36} - \frac{84}{36} \right) = 0$$

$$3 \left(\left(x - \frac{5}{6}\right)^2 - \frac{109}{36} \right) = 0$$

We next apply the difference of squares theorem. $\frac{109}{36}$ is the square of a number - its own square root.

$$\frac{109}{36} = \left(\sqrt{\frac{109}{36}} \right)^2 = \left(\frac{\sqrt{109}}{\sqrt{36}} \right)^2 = \left(\frac{\sqrt{109}}{6} \right)^2$$

$$3 \left(\left(x - \frac{5}{6} \right)^2 - \left(\frac{\sqrt{109}}{6} \right)^2 \right) = 0$$

$$3 \left(x - \frac{5}{6} + \frac{\sqrt{109}}{6} \right) \left(x - \frac{5}{6} - \frac{\sqrt{109}}{6} \right) = 0$$

We solve for the zeroes of both linear factors:

$$x - \frac{5}{6} + \frac{\sqrt{109}}{6} = 0 \quad \text{or} \quad x - \frac{5}{6} - \frac{\sqrt{109}}{6} = 0$$

$$x + \frac{\sqrt{109}}{6} = \frac{5}{6} \quad \quad \quad x - \frac{\sqrt{109}}{6} = \frac{5}{6}$$

$$x_1 = \frac{5}{6} - \frac{\sqrt{109}}{6} = \boxed{\frac{5 - \sqrt{109}}{6}} \quad \quad \quad x_2 = \frac{5}{6} + \frac{\sqrt{109}}{6} = \boxed{\frac{5 + \sqrt{109}}{6}}$$

We check: if $x = \frac{5 - \sqrt{109}}{6}$, then

$$\begin{aligned} \text{LHS} &= 3x^2 - 5x - 7 = 3 \left(\frac{5 - \sqrt{109}}{6} \right)^2 - 5 \left(\frac{5 - \sqrt{109}}{6} \right) - 7 \\ &= \frac{3}{1} \cdot \frac{(5 - \sqrt{109})^2}{36} - \frac{5}{1} \left(\frac{5 - \sqrt{109}}{6} \right) - 7 = \\ &= \frac{3(25 - 5\sqrt{109} - 5\sqrt{109} + 109)}{36} - \frac{5(5 - \sqrt{109})}{6} - 7 = \frac{134 - 10\sqrt{109}}{12} - \frac{25 - 5\sqrt{109}}{6} - 7 \\ &= \frac{134 - 10\sqrt{109}}{12} - \frac{2(25 - 5\sqrt{109})}{12} - 7 = \frac{134 - 10\sqrt{109} - 2(25 - 5\sqrt{109})}{12} - 7 \\ &= \frac{134 - 10\sqrt{109} - 50 + 10\sqrt{109}}{12} - 7 = \frac{84}{12} - 7 = 7 - 7 = 0 = \text{RHS} \end{aligned}$$

and if $x = \frac{5 + \sqrt{109}}{6}$, then

$$\begin{aligned} \text{LHS} &= 3x^2 - 5x - 7 = 3 \left(\frac{5 + \sqrt{109}}{6} \right)^2 - 5 \left(\frac{5 + \sqrt{109}}{6} \right) - 7 \\ &= \frac{3}{1} \cdot \frac{(5 + \sqrt{109})^2}{36} - \frac{5}{1} \left(\frac{5 + \sqrt{109}}{6} \right) - 7 \\ &= \frac{3(25 + 5\sqrt{109} + 5\sqrt{109} + 109)}{36} - \frac{5(5 + \sqrt{109})}{6} - 7 = \frac{134 + 10\sqrt{109}}{12} - \frac{25 + 5\sqrt{109}}{6} - 7 \\ &= \frac{134 + 10\sqrt{109}}{12} - \frac{2(25 + 5\sqrt{109})}{12} - 7 = \frac{134 + 10\sqrt{109} - 2(25 + 5\sqrt{109})}{12} - 7 \\ &= \frac{134 + 10\sqrt{109} - 50 - 10\sqrt{109}}{12} - 7 = \frac{84}{12} - 7 = 7 - 7 = 0 = \text{RHS} \end{aligned}$$

Thus our solution, $\frac{5 + \sqrt{109}}{6}$ and $\frac{5 - \sqrt{109}}{6}$ is correct.

2. Find all numbers that are exactly three less than their own square.

Solution: Let us denote this number by x . Then its square is x^2 and

$$\begin{aligned}
 x &= x^2 - 3 && \text{reduce one side to zero} \\
 0 &= x^2 - x - 3 \\
 x^2 - x - 3 &= 0 && \left(x - \frac{1}{2}\right)^2 = x^2 - x + \frac{1}{4} \\
 \underbrace{x^2 - x + \frac{1}{4}} - \frac{1}{4} - 3 &= 0 && 3 = \frac{12}{4} \\
 \left(x - \frac{1}{2}\right)^2 - \frac{13}{4} &= 0 && \sqrt{\frac{13}{4}} = \frac{\sqrt{13}}{\sqrt{4}} = \frac{\sqrt{13}}{2} \\
 \left(x - \frac{1}{2}\right)^2 - \left(\frac{\sqrt{13}}{2}\right)^2 &= 0 \\
 \left(x - \frac{1}{2} + \frac{\sqrt{13}}{2}\right) \left(x - \frac{1}{2} - \frac{\sqrt{13}}{2}\right) &= 0
 \end{aligned}$$

$$\begin{aligned}
 \text{Either } x - \frac{1}{2} + \frac{\sqrt{13}}{2} &= 0 && \text{or} && x - \frac{1}{2} - \frac{\sqrt{13}}{2} = 0 \\
 x + \frac{\sqrt{13}}{2} &= \frac{1}{2} && && x - \frac{\sqrt{13}}{2} = \frac{1}{2} \\
 x_1 &= \frac{1}{2} - \frac{\sqrt{13}}{2} = \frac{1 - \sqrt{13}}{2} && && x_2 = \frac{1}{2} + \frac{\sqrt{13}}{2} = \frac{1 + \sqrt{13}}{2}
 \end{aligned}$$

We check: If $x_1 = \frac{1 - \sqrt{13}}{2}$, then

$$\begin{aligned}
 x_1^2 - 3 &= \left(\frac{1 - \sqrt{13}}{2}\right)^2 - 3 = \frac{(1 - \sqrt{13})^2}{2^2} - 3 = \frac{1 + 13 - 2\sqrt{13}}{4} - 3 = \frac{14 - 2\sqrt{13}}{4} - 3 = \frac{2(7 - \sqrt{13})}{4} - 3 \\
 &= \frac{7 - \sqrt{13}}{2} - \frac{6}{2} = \frac{1 - \sqrt{13}}{2} = x_1 \checkmark \quad \text{Thus } x_1 = \frac{1 - \sqrt{13}}{2} \text{ works.}
 \end{aligned}$$

Let us see about the other number. If $x_2 = \frac{1 + \sqrt{13}}{2}$, then

$$\begin{aligned}
 x_2^2 - 3 &= \left(\frac{1 + \sqrt{13}}{2}\right)^2 - 3 = \frac{(1 + \sqrt{13})^2}{2^2} - 3 = \frac{1 + 13 + 2\sqrt{13}}{4} - 3 = \frac{14 + 2\sqrt{13}}{4} - 3 = \frac{2(7 + \sqrt{13})}{4} - 3 \\
 &= \frac{7 + \sqrt{13}}{2} - \frac{6}{2} = \frac{1 + \sqrt{13}}{2} = x_2 \checkmark \quad \text{Thus } x_2 = \frac{1 + \sqrt{13}}{2} \text{ also works.}
 \end{aligned}$$

We found two numbers; they both work.