

Complex fractions are fractions with more than one fraction bars. For example, $\frac{\frac{1}{2} - 3}{5 + \frac{1}{4}}$ and $\frac{1 - \frac{2}{x+1}}{\frac{5}{x-3} + \frac{x}{2}}$ are complex fractions.

A complex fraction can always be simplified, that is, brought to a form with only one fraction bar. We will present two methods. The first one might be convenient because it is familiar and consists of steps we know very well. The other method is new to us. However, it appears to be a very effective method.

Example 1. Simplify the given complex fractions.

$$\text{a) } \frac{\frac{2}{3} - \frac{1}{4}}{\frac{3}{8} + \frac{1}{2}} \quad \text{b) } \frac{\frac{2}{x} - 5}{1 + \frac{3}{x}}$$

Solution 1: a) This method involves performing operations on fractions: a subtraction (in the numerator), an addition (in the denominator), and finally a division.

$$\text{The subtraction: } \frac{2}{3} - \frac{1}{4} = \frac{8}{12} - \frac{3}{12} = \frac{5}{12}$$

$$\text{The addition: } \frac{3}{8} + \frac{1}{2} = \frac{3}{8} + \frac{4}{8} = \frac{7}{8}$$

$$\text{The division: } \frac{\frac{5}{12}}{\frac{7}{8}} = \frac{5}{12} \cdot \frac{8}{7} = \frac{5 \cdot 2 \cdot \cancel{4}}{3 \cdot \cancel{4} \cdot 7} = \boxed{\frac{10}{21}}$$

We can perform these operations in a single computation:

$$\frac{\frac{2}{3} - \frac{1}{4}}{\frac{3}{8} + \frac{1}{2}} = \frac{\frac{8-3}{12}}{\frac{3+4}{8}} = \frac{\frac{5}{12}}{\frac{7}{8}} = \frac{5}{12} \cdot \frac{8}{7} = \frac{40}{84} = \boxed{\frac{10}{21}}$$

b) We can apply the same steps with the fraction containing variables.

$$\text{The subtraction in the numerator: } \frac{2}{x} - 5 = \frac{2}{x} - \frac{5}{1} = \frac{2}{x} - \frac{5x}{x} = \frac{2-5x}{x} = \frac{-5x+2}{x}$$

$$\text{The addition in the denominator: } 1 + \frac{3}{x} = \frac{x}{x} + \frac{3}{x} = \frac{x+3}{x}$$

$$\text{The division: } \frac{\frac{-5x+2}{x}}{\frac{x+3}{x}} = \frac{-5x+2}{x} \cdot \frac{x}{x+3} = \frac{-5x+2}{\cancel{x}} \cdot \frac{\cancel{x}}{x+3} = \boxed{\frac{-5x+2}{x+3}}$$

Solution 2: a) This method involves the fundamental property of fractions. Recall that we do not change the value of a fraction when multiplying both numerator and denominator by the same non-zero number. This technique involves 'blasting' the denominators.

Consider the fraction $\frac{\frac{2}{3} - \frac{1}{4}}{\frac{3}{8} + \frac{1}{2}}$. If we multiplied the numerator, $\frac{2}{3} - \frac{1}{4}$ by 12, that would clear the denominator

of both $\frac{2}{3}$ and $\frac{1}{4}$. If we multiplied the denominator, $\frac{3}{8} + \frac{1}{2}$ by 8, that would clear both denominators. Since we must use the same number for both numerator and denominator, we will multiply both by 24. This technique is surprisingly powerful: we get the final result much faster than before.

$$\frac{\frac{2}{3} - \frac{1}{4}}{\frac{3}{8} + \frac{1}{2}} = \frac{24 \left(\frac{2}{3} - \frac{1}{4} \right)}{24 \left(\frac{3}{8} + \frac{1}{2} \right)} = \frac{24 \cdot \frac{2}{3} - 24 \cdot \frac{1}{4}}{24 \cdot \frac{3}{8} + 24 \cdot \frac{1}{2}} = \frac{\overset{8}{\cancel{24}} \cdot \frac{2}{\cancel{3}} - \overset{6}{\cancel{24}} \cdot \frac{1}{\cancel{4}}}{\overset{3}{\cancel{24}} \cdot \frac{3}{\cancel{8}} + \overset{12}{\cancel{24}} \cdot \frac{1}{\cancel{2}}} = \frac{8 \cdot 2 - 6 \cdot 1}{3 \cdot 3 + 12 \cdot 1} = \frac{16 - 6}{9 + 12} = \boxed{\frac{10}{21}}$$

b) We can apply the same steps with the fraction containing variables. In the case of $\frac{\frac{2}{x} - 5}{1 + \frac{3}{x}}$, both numerator and denominator would be cleared when multiplied by x .

$$\frac{\frac{2}{x} - 5}{1 + \frac{3}{x}} = \frac{x \left(\frac{2}{x} - 5 \right)}{x \left(1 + \frac{3}{x} \right)} = \frac{x \cdot \frac{2}{x} - x \cdot 5}{x \cdot 1 + x \cdot \frac{3}{x}} = \frac{\cancel{x} \cdot \frac{2}{\cancel{x}} - x \cdot 5}{x \cdot 1 + \cancel{x} \cdot \frac{3}{\cancel{x}}} = \frac{2 - 5x}{x + 3} = \boxed{\frac{-5x + 2}{x + 3}}$$

There might be cancelations after we simplify the complex fractions. Also, sometimes complex fractions are given using negative exponents.

Example 2. Simplify the complex fraction $\frac{25 - x^{-2}}{5 + x^{-1}}$

Solution 1: We first re-write the expression without negative exponents.

$$\begin{aligned} \frac{25 - x^{-2}}{5 + x^{-1}} &= \frac{25 - \frac{1}{x^2}}{5 + \frac{1}{x}} = \frac{\frac{25x^2}{x^2} - \frac{1}{x^2}}{\frac{5x}{x} + \frac{1}{x}} = \frac{\frac{25x^2 - 1}{x^2}}{\frac{5x + 1}{x}} = \frac{(5x + 1)(5x - 1)}{x^2} \cdot \frac{x}{5x + 1} \\ &= \frac{\cancel{(5x + 1)}(5x - 1)}{\cancel{x}x} \cdot \frac{\cancel{x}}{\cancel{5x + 1}} = \boxed{\frac{5x - 1}{x}} \end{aligned}$$



Sample Problems

Simplify each of the following expressions.

$$1. \frac{\frac{1}{2} + \frac{1}{3}}{\frac{1}{2} - \frac{1}{3}}$$

$$3. \frac{1 - \frac{1}{x^2}}{1 + \frac{1}{x}}$$

$$5. \frac{\frac{3}{x-1} - 1}{\frac{2}{x-1} + 1}$$

$$8. \frac{a^{-1} - b^{-1}}{a^{-2} - b^{-2}}$$

$$2. \frac{5 - \frac{1}{a}}{\frac{1}{a^2} - 25}$$

$$4. \frac{\frac{4}{a^2} - 1}{1 - \frac{2}{a}}$$

$$6. \frac{2 - \frac{3}{x+1}}{3 - \frac{2x}{x+1}}$$

$$9. \frac{x^2 - 2y^{-3}}{x^{-1} + 3y^2}$$

$$7. 1 - \frac{1}{1 - \frac{1}{x-3}}$$

$$10. (x - y^{-2})^{-3}$$



Practice Problems

Simplify each of the following expressions.

$$1. \frac{2a - \frac{1}{8a}}{4 + \frac{1}{a}}$$

$$6. \frac{\frac{1}{x+a} + \frac{1}{x-a}}{\frac{1}{x+a} - \frac{1}{x-a}}$$

$$10. \frac{1 - \frac{y}{y-1}}{\frac{y}{y+1} - 1}$$

$$14. \frac{1 - \frac{2}{3 - \frac{4}{p}}}{1 + \frac{2}{3 - \frac{4}{p}}}$$

$$2. \frac{1}{2 - \frac{1}{m-4}}$$

$$7. 1 - \frac{1}{1 - \frac{1}{1 - \frac{1}{q-1}}}$$

$$11. \frac{\frac{4}{z-6} - \frac{7}{z}}{\frac{5}{z} + \frac{7}{z-6}}$$

$$15. \frac{9a^{-2} - b^{-2}}{3a^{-1} + b^{-1}}$$

$$3. \frac{1}{1 - \frac{1}{x-1}}$$

$$8. \frac{6 + \frac{2}{x}}{3x + 1}$$

$$12. \frac{\frac{4}{x^2} - \frac{3}{x}}{\frac{1}{x^2} + \frac{2}{3x}}$$

$$16. \frac{x^{-3}y - 2x^2y^{-1}}{x^{-1} + y^{-1}}$$

$$4. \frac{\frac{1}{a} + \frac{1}{b}}{\frac{1}{a} - \frac{1}{b}}$$

$$9. \frac{2}{\frac{1}{a} + \frac{1}{b}}$$

$$13. \frac{\frac{5}{x-3} - \frac{1}{x}}{\frac{2}{x} + \frac{3}{x-3}}$$

$$17. (3 - a^{-2})^{-2}$$

$$5. \frac{5 - \frac{x}{x-1}}{2 - \frac{x}{x+1}}$$



Answers


Sample Problems

$$\begin{array}{lllllll}
 1. 5 & 2. -\frac{a}{5a+1} & 3. \frac{x-1}{x} & 4. -\frac{a+2}{a} & 5. \frac{-x+4}{x+1} & 6. \frac{2x-1}{x+3} & 7. -\frac{1}{x-4} \\
 8. \frac{ab}{a+b} & 9. \frac{x^3y^3-2x}{y^3+3xy^5} \text{ or } \frac{x(x^2y^3-2)}{y^3(1+3xy^2)} & & & 10. \frac{y^6}{(xy^2-1)^3} & &
 \end{array}$$

Practice Problems

$$\begin{array}{llllllll}
 1. \frac{4a-1}{8} & 2. \frac{m-4}{2m-9} & 3. \frac{x-1}{x-2} & 4. \frac{a+b}{b-a} & 5. \frac{4x^2-x-5}{x^2+x-2} & 6. -\frac{x}{a} & 7. q-1 & 8. \frac{8}{x} \\
 9. \frac{2ab}{a+b} & 10. \frac{y+1}{y-1} & 11. \frac{14-z}{4z-10} \text{ or } \frac{14-z}{2(2z-5)} & 12. \frac{12-9x}{2x+3} \text{ or } \frac{-3(3x-4)}{2x+3} & 13. \frac{4x+3}{5x-6} \\
 14. \frac{p-4}{5p-4} & 15. \frac{3b-a}{ab} & 16. \frac{y^2-2x^5}{x^3+x^2y} \text{ or } \frac{y^2-2x^5}{x^2(x+y)} & 17. \frac{a^4}{(1-3a^2)^2}
 \end{array}$$

Sample Problems



Solutions

Simplify each of the following expressions.

$$1. \frac{\frac{1}{2} + \frac{1}{3}}{\frac{1}{2} - \frac{1}{3}}$$

Solution: We will first perform the addition indicated in the numerator and denominator. After the addition and subtraction, we will divide by multiplying by the reciprocal.

$$\frac{\frac{1}{2} + \frac{1}{3}}{\frac{1}{2} - \frac{1}{3}} = \frac{\frac{1 \cdot 3}{2 \cdot 3} + \frac{1 \cdot 2}{3 \cdot 2}}{\frac{1 \cdot 3}{2 \cdot 3} - \frac{1 \cdot 2}{3 \cdot 2}} = \frac{\frac{3}{6} + \frac{2}{6}}{\frac{3}{6} - \frac{2}{6}} = \frac{\frac{2+3}{6}}{\frac{3-2}{6}} = \frac{\frac{5}{6}}{\frac{1}{6}} = \frac{5}{\cancel{6}} \cdot \frac{\cancel{6}}{1} = \frac{5}{1} = \boxed{5}$$

$$2. \frac{5 - \frac{1}{a}}{\frac{1}{a^2} - 25}$$

Solution: Method 1. Start with the subtractions. We need to work with common denominators. The numerator:

$$5 - \frac{1}{a} = \frac{5}{1} - \frac{1}{a} = \frac{5a}{a} - \frac{1}{a} = \frac{5a - 1}{a}$$

The denominator:

$$\frac{1}{a^2} - 25 = \frac{1}{a^2} - \frac{25}{1} = \frac{1}{a^2} - \frac{25a^2}{a^2} = \frac{1 - 25a^2}{a^2}$$

Notice that the numerator here factors via the difference of squares theorem.

$$\frac{1 - 25a^2}{a^2} = \frac{1^2 - (5a)^2}{a^2} = \frac{(1 - 5a)(1 + 5a)}{a^2}$$

Now we are ready to perform the division. To divide is to multiply by the reciprocal.

$$\begin{aligned} \frac{5 - \frac{1}{a}}{\frac{1}{a^2} - 25} &= \frac{\frac{5a - 1}{a}}{\frac{(1 - 5a)(1 + 5a)}{a^2}} = \frac{5a - 1}{a} \cdot \frac{a^2}{(1 - 5a)(1 + 5a)} \quad \text{cancel out } a \\ &= \frac{5a - 1}{\cancel{a}} \cdot \frac{\cancel{a^2}}{(1 - 5a)(1 + 5a)} = \frac{a(5a - 1)}{(1 - 5a)(1 + 5a)} \end{aligned}$$

There is one more cancellation: $1 - 5a$ and $5a - 1$ are opposites. We re-write $5a - 1$ as $-(1 - 5a)$. Then

$$\frac{a(5a - 1)}{(1 - 5a)(1 + 5a)} = \frac{-(1 - 5a)a}{(1 - 5a)(1 + 5a)} = \frac{\cancel{-(1 - 5a)}a}{\cancel{(1 - 5a)}(1 + 5a)} = \boxed{\frac{-a}{1 + 5a} \quad \text{or} \quad -\frac{a}{5a + 1}}$$

Method 2. This method is very efficient. The main idea is to clear denominators by multiplying both numerator and denominator by the same quantity. We usually do not apply this method with usual fractions but it does work there too. The first step is to multiply both numerator and denominator by a^2 , the quantity that clears the denominator in the fractions in both numerator and denominator.

$$\frac{5 - \frac{1}{a}}{\frac{1}{a^2} - 25} = \frac{5 - \frac{1}{a}}{\frac{1}{a^2} - 25} \cdot \frac{a^2}{a^2} = \frac{a^2 \left(5 - \frac{1}{a}\right)}{a^2 \left(\frac{1}{a^2} - 25\right)} = \frac{5a^2 - a^2 \cdot \frac{1}{a}}{a^2 \cdot \frac{1}{a^2} - 25a^2} = \frac{5a^2 - a}{1 - 25a^2}$$

We now factor the expressions in the numerator and denominator so that we could see if further simplification is possible. We factor out the greatest common factor from the numerator and factor the denominator via the difference of squares theorem.

$$\frac{5a^2 - a}{1 - 25a^2} = \frac{a(5a - 1)}{(1 - 5a)(1 + 5a)}$$

Notice that $5a - 1$ and $1 - 5a$ are opposites. We factor out -1 from the numerator, so we re-write $5a - 1$ as $-(1 - 5a)$. Then we cancel out the common factor.

$$\frac{a(5a - 1)}{(1 - 5a)(1 + 5a)} = \frac{a(-1)(1 - 5a)}{(1 - 5a)(1 + 5a)} = \frac{a(-1)\cancel{(1 - 5a)}}{\cancel{(1 - 5a)}(1 + 5a)} = \frac{a(-1)}{1 + 5a} = \boxed{\frac{-a}{5a + 1}}$$

$$3. \frac{1 - \frac{1}{x^2}}{1 + \frac{1}{x}}$$

Solution: Method 1. We bring fractions to the common denominator and add and subtract. Then we perform the division as multiplication by the reciprocal. Finally, we factor and cancel out common factors.

$$\frac{1 - \frac{1}{x^2}}{1 + \frac{1}{x}} = \frac{\frac{1 \cdot x^2}{1 \cdot x^2} - \frac{1}{x^2}}{\frac{1 \cdot x}{1 \cdot x} + \frac{1}{x}} = \frac{\frac{x^2 - 1}{x^2}}{\frac{x + 1}{x}} = \frac{x^2 - 1}{x^2} \cdot \frac{x}{x + 1} = \frac{(x + 1)(x - 1)}{x^2} \cdot \frac{x}{x + 1} = \frac{\cancel{(x + 1)}(x - 1)}{x^{\cancel{2}} \cdot \cancel{x + 1}} = \boxed{\frac{x - 1}{x}}$$

Method 2. First we multiply both numerator and denominator by x^2 .

$$\frac{1 - \frac{1}{x^2}}{1 + \frac{1}{x}} = \frac{1 - \frac{1}{x^2}}{1 + \frac{1}{x}} \cdot \frac{x^2}{x^2} = \frac{x^2 \left(1 - \frac{1}{x^2}\right)}{x^2 \left(1 + \frac{1}{x}\right)} = \frac{x^2 \cdot 1 - x^2 \cdot \frac{1}{x^2}}{x^2 \cdot 1 + x^2 \cdot \frac{1}{x}} = \frac{x^2 - 1}{x^2 + x}$$

We now factor and cancel out common factors.

$$\frac{x^2 - 1}{x^2 + x} = \frac{(x + 1)(x - 1)}{x(x + 1)} = \frac{\cancel{(x + 1)}(x - 1)}{x\cancel{(x + 1)}} = \boxed{\frac{x - 1}{x}}$$

$$4. \frac{\frac{4}{a^2} - 1}{1 - \frac{2}{a}}$$

Solution: Method 1

$$\begin{aligned} \frac{\frac{4}{a^2} - 1}{1 - \frac{2}{a}} &= \frac{\frac{4}{1} - \frac{1}{1}}{\frac{1}{1} - \frac{2}{a}} = \frac{\frac{4}{1} - \frac{1 \cdot a^2}{1 \cdot a^2}}{\frac{1 \cdot a}{1 \cdot a} - \frac{2}{a}} = \frac{\frac{4 - a^2}{a^2}}{\frac{a - 2}{a}} = \frac{4 - a^2}{a^2} \cdot \frac{a}{a - 2} = \frac{(2 + a)(2 - a)}{a} \cdot \frac{1}{a - 2} \\ &= \frac{(2 + a)(-1)(a - 2)}{a(a - 2)} = \frac{(2 + a)(-1)\cancel{(a - 2)}}{a\cancel{(a - 2)}} = \boxed{\frac{-a + 2}{a}} \end{aligned}$$

Method 2. We first multiply both numerator and denominator by a^2 . Then we factor all expressions and simplify the fraction.

$$\frac{\frac{4}{a^2} - 1}{1 - \frac{2}{a}} = \frac{a^2 \left(\frac{4}{a^2} - 1 \right)}{a^2 \left(1 - \frac{2}{a} \right)} = \frac{4 - a^2}{a^2 - 2a} = \frac{(2 + a)(2 - a)}{a(a - 2)}$$

Notice that $a - 2$ and $2 - a$ are opposites. We will factor out -1 from $2 - a$ in the numerator.

$$\frac{(2 + a)(2 - a)}{a(a - 2)} = \frac{(2 + a)(-1)(a - 2)}{a(a - 2)} = \frac{-(2 + a)\cancel{(a - 2)}}{a\cancel{(a - 2)}} = \frac{-(2 + a)}{a} = \boxed{\frac{-a + 2}{a}}$$

$$5. \frac{\frac{3}{x-1} - 1}{\frac{2}{x-1} + 1}$$

Solution: Method 1.

$$\begin{aligned} \frac{\frac{3}{x-1} - 1}{\frac{2}{x-1} + 1} &= \frac{\frac{3}{x-1} - \frac{1}{1}}{\frac{2}{x-1} + \frac{1}{1}} = \frac{\frac{3}{x-1} - \frac{1 \cdot (x-1)}{1 \cdot (x-1)}}{\frac{2}{x-1} + \frac{1 \cdot (x-1)}{1 \cdot (x-1)}} = \frac{\frac{3}{x-1} - \frac{x-1}{x-1}}{\frac{2}{x-1} + \frac{x-1}{x-1}} = \frac{\frac{3 - (x-1)}{x-1}}{\frac{2 + (x-1)}{x-1}} = \frac{3 - x + 1}{2 + x - 1} = \frac{3 - x + 1}{x - 1} \\ &= \frac{-x + 4}{x - 1} = \frac{-x + 4}{x - 1} \cdot \frac{x + 1}{x + 1} = \frac{-x + 4}{\cancel{x - 1}} \cdot \frac{\cancel{x - 1}}{x + 1} = \boxed{\frac{-x + 4}{x + 1}} \end{aligned}$$

Method 2. We multiply both numerator and denominator by $x - 1$. We need to be careful with subtractions as we are subtracting entire expressions.

$$\frac{\frac{3}{x-1} - 1}{\frac{2}{x-1} + 1} = \frac{(x-1) \left(\frac{3}{x-1} - 1 \right)}{(x-1) \left(\frac{2}{x-1} + 1 \right)} = \frac{3 - 1(x-1)}{2 + 1(x-1)} = \frac{3 - x + 1}{2 + x - 1} = \frac{4 - x}{x + 1} = \boxed{\frac{-x + 4}{x + 1}}$$

$$6. \frac{2 - \frac{3}{x+1}}{3 - \frac{2x}{x+1}}$$

Solution: Method 1. We first work out the numerator

$$\begin{aligned} 2 - \frac{3}{x+1} &= \frac{2}{1} - \frac{3}{x+1} \quad \text{common denominator is } x+1 \\ &= \frac{2(x+1)}{x+1} - \frac{3}{x+1} = \frac{2(x+1) - 3}{x+1} = \frac{2x+2-3}{x+1} = \frac{2x-1}{x+1} \end{aligned}$$

Now for the denominator

$$\begin{aligned} 3 - \frac{2x}{x+1} &= \frac{3}{1} - \frac{2x}{x+1} \quad \text{common denominator is } x+1 \\ &= \frac{3(x+1)}{x+1} - \frac{2x}{x+1} = \frac{3(x+1) - 2x}{x+1} = \frac{3x+3-2x}{x+1} = \frac{x+3}{x+1} \end{aligned}$$

Now the division:

$$\begin{aligned} \frac{2 - \frac{3}{x+1}}{3 - \frac{2x}{x+1}} &= \frac{\frac{2x-1}{x+1}}{\frac{x+3}{x+1}} = \quad \text{to divide is to multiply by the reciprocal} \\ &= \frac{2x-1}{x+1} \cdot \frac{x+1}{x+3} = \frac{2x-1}{x+3} \cdot \frac{\cancel{x+1}}{\cancel{x+1}} = \boxed{\frac{2x-1}{x+3}} \end{aligned}$$

Method 2.

$$\frac{2 - \frac{3}{x+1}}{3 - \frac{2x}{x+1}} \cdot \frac{x+1}{x+1} = \frac{\left(2 - \frac{3}{x+1}\right)(x+1)}{\left(3 - \frac{2x}{x+1}\right)(x+1)} = \frac{2(x+1) - \frac{3}{x+1}(x+1)}{3(x+1) - \frac{2x}{x+1}(x+1)} = \frac{2(x+1) - 3}{3(x+1) - 2x} = \frac{2x+2-3}{3x+3-2x} = \boxed{\frac{2x-1}{x+3}}$$

$$7. 1 - \frac{1}{1 - \frac{1}{x-3}}$$

Solution:

$$\begin{aligned} 1 - \frac{1}{1 - \frac{1}{x-3}} &= 1 - \frac{1}{\frac{1}{1} - \frac{1}{x-3}} = 1 - \frac{1}{\frac{1 \cdot (x-3)}{1 \cdot (x-3)} - \frac{1}{x-3}} = 1 - \frac{1}{\frac{x-3}{x-3} - \frac{1}{x-3}} \\ &= 1 - \frac{1}{\frac{x-3-1}{x-3}} = 1 - \frac{1}{\frac{x-4}{x-3}} = 1 - \frac{1}{1} \cdot \frac{x-3}{x-4} = 1 - \frac{x-3}{x-4} \\ &= \frac{x-4}{x-4} - \frac{x-3}{x-4} = \frac{(x-4) - (x-3)}{x-4} = \frac{x-4-x+3}{x-4} = \boxed{\frac{-1}{x-4}} \end{aligned}$$

$$8. \frac{a^{-1} - b^{-1}}{a^{-2} - b^{-2}}$$

Solution: This does not even look like a complex fraction, but it will, once we re-write it positive exponents. Recall that $a^{-n} = \frac{1}{a^n}$. So our problem can be re-written as

$$\frac{a^{-1} - b^{-1}}{a^{-2} - b^{-2}} = \frac{\frac{1}{a} - \frac{1}{b}}{\frac{1}{a^2} - \frac{1}{b^2}}$$

Method 1.

$$\frac{\frac{1}{a} - \frac{1}{b}}{\frac{1}{a^2} - \frac{1}{b^2}} = \frac{\frac{b-a}{ab}}{\frac{b^2-a^2}{a^2b^2}} = \frac{b-a}{ab} \cdot \frac{a^2b^2}{b^2-a^2} = \frac{(b-a)ab}{b^2-a^2} = \frac{(b-a)ab}{(b-a)(b+a)} = \frac{\cancel{(b-a)}ab}{\cancel{(b-a)}(b+a)} = \frac{ab}{b+a} = \boxed{\frac{ab}{a+b}}$$

Method 2.

$$\frac{\frac{1}{a} - \frac{1}{b}}{\frac{1}{a^2} - \frac{1}{b^2}} = \frac{\frac{1}{a} - \frac{1}{b}}{\frac{1}{a^2} - \frac{1}{b^2}} \cdot \frac{a^2b^2}{a^2b^2} = \frac{a^2b^2 \left(\frac{1}{a} - \frac{1}{b} \right)}{a^2b^2 \left(\frac{1}{a^2} - \frac{1}{b^2} \right)} = \frac{ab^2 - a^2b}{b^2 - a^2} = \frac{ab\cancel{(b-a)}}{\cancel{(b-a)}(b+a)} = \frac{ab(b-a)}{(b-a)(b+a)} = \boxed{\frac{ab}{a+b}}$$

$$9. \frac{x^2 - 2y^{-3}}{x^{-1} + 3y^2}$$

Solution: Method 1.

$$\frac{x^2 - 2y^{-3}}{x^{-1} + 3y^2} = \frac{x^2 - 2\frac{1}{y^3}}{\frac{1}{x} + 3y^2} = \frac{\frac{x^2}{1} - \frac{2}{y^3}}{\frac{1}{x} + \frac{3y^2}{1}} = \frac{\frac{x^2y^3}{y^3} - \frac{2}{y^3}}{\frac{1}{x} + \frac{3xy^2}{x}} = \frac{\frac{x^2y^3 - 2}{y^3}}{\frac{1 + 3xy^2}{x}} = \frac{x^2y^3 - 2}{y^3} \cdot \frac{x}{1 + 3xy^2} = \boxed{\frac{x(x^2y^3 - 2)}{y^3(1 + 3xy^2)}}$$

Method 2.

$$\frac{x^2 - 2y^{-3}}{x^{-1} + 3y^2} = \frac{x^2 - 2\frac{1}{y^3}}{\frac{1}{x} + 3y^2} = \frac{\frac{x^2}{1} - \frac{2}{y^3}}{\frac{1}{x} + \frac{3y^2}{1}} \cdot \frac{xy^3}{xy^3} = \frac{xy^3 \left(\frac{x^2}{1} - \frac{2}{y^3} \right)}{xy^3 \left(\frac{1}{x} + \frac{3y^2}{1} \right)} = \frac{x^3y^3 - 2x}{y^3 + 3xy^5} = \boxed{\frac{x(x^2y^3 - 2)}{y^3(1 + 3xy^2)}}$$

$$10. (x - y^{-2})^{-3}$$

Solution: Method 1.

$$\begin{aligned} (x - y^{-2})^{-3} &= \frac{1}{(x - y^{-2})^3} = \frac{1}{\left(x - \frac{1}{y^2}\right)^3} = \frac{1}{\left(\frac{xy^2}{y^2} - \frac{1}{y^2}\right)^3} = \frac{1}{\left(\frac{xy^2 - 1}{y^2}\right)^3} = \frac{1}{\frac{(xy^2 - 1)^3}{(y^2)^3}} \\ &= 1 \cdot \frac{(y^2)^3}{(xy^2 - 1)^3} = \boxed{\frac{y^6}{(xy^2 - 1)^3}} \end{aligned}$$

For more documents like this, visit our page at <https://teaching.martahidegkuti.com> and click on Lecture Notes. E-mail questions or comments to mhidegkuti@ccc.edu.