

Practice Problems

1. Differentiate each of the following. Make sure to simplify your final answer.

a) $f(x) = 5^{-2x+1}$

g) $L(\theta) = \frac{\tan \theta - \cot \theta}{\tan \theta}$

m) $g(w) = \cos^{-1}(w^2 + 1)$

b) $m(y) = \log_7(\sqrt{y^6 + 4y^2 + 10})$

h) $v(t) = e^{2t} - e^{-2t}$

n) $f(y) = 2^{\log_3 y}$

c) $f(m) = \frac{1}{3}m^3 \ln m - \frac{1}{9}m^3$

i) $f(x) = \log_2(x^{10})$

o) $f(\theta) = \cot(5\theta)$

d) $m(p) = \sqrt{1 - p^2}$

j) $g(x) = \tan^{-1}(5x)$

p) $g(x) = \sqrt{x + \sqrt{x}}$

e) $g(t) = e^{\sin t - \cos t}$

k) $h(\alpha) = \log_2(\tan 2\alpha)$

q) $f(\beta) = \csc(\beta^2 + 1)$

f) $f(a) = \frac{2^{3a-1}}{2^{-a+1}}$

l) $f(x) = \cos(\sin^{-1} x)$

r) $m(y) = \sec^{-1}(3y)$

s) $P(t) = \log_5(\csc t)$

2. Compute each of the following integrals.

a) $\int \frac{1}{(2a-7)^4} da$

d) $\int_0^1 \frac{1}{1+\theta^2} d\theta$

g) $\int \frac{1}{|x|\sqrt{x^2-1}} dx$

b) $\int \frac{1}{2a-7} da$

e) $\int \frac{1}{\sqrt{1-y^2}} dy$

h) $\int \csc x \cot x dx$

c) $\int_0^2 e^{-2t+1} dt$

f) $\int_0^3 2^x dx$

i) $\int_0^{\pi/4} \tan^2 x dx$

3. Use the Fundamental Theorem of Calculus to compute each of the following derivatives.

a) $\frac{d}{dx} \left(\int_1^x \sqrt{t} dt \right)$

c) $\frac{d}{dx} \left(\int_1^{9x} \sqrt{t} dt \right)$

e) $\frac{d}{dx} \left(\int_0^{x^2} \frac{1}{1+t^2} dt \right)$

b) $\frac{d}{dx} \left(\int_x^{16} \sqrt{t} dt \right)$

d) $\frac{d}{dx} \left(\int_0^x \frac{1}{1+t^2} dt \right)$

f) $\frac{d}{dx} \left[\left(\int_0^x \frac{1}{1+t^2} dt \right)^2 \right]$

Practice Problems - Answers

1.) a) $f'(x) = -2(\ln 5) 5^{-2x+1}$

b) $m'(y) = \frac{3y^5 + 4y}{\ln 7 (y^6 + 4y^2 + 10)}$

c) $f'(m) = m^2 \ln m$

d) $m'(p) = -\frac{p}{\sqrt{1-p^2}}$

e) $g'(t) = (\cos t + \sin t) e^{\sin t - \cos t}$

f) $f'(a) = (\ln 2) 2^{4a}$

$$\text{g) } L'(\theta) = \frac{2 \cos \theta}{\sin^3 \theta} = 2 \cot^3 \theta + 2 \cot \theta \quad \text{h) } v'(t) = 2e^{2t} + 2e^{-2t} \quad \text{i) } f'(x) = \frac{10}{x \ln 2}$$

$$\text{j) } g'(x) = \frac{5}{25x^2 + 1} \quad \text{k) } h'(\alpha) = \frac{2}{\ln 2} (\cot 2\alpha + \tan 2\alpha) = \frac{2 \tan^2 2\alpha + 2}{(\ln 2) \tan 2\alpha} \quad \text{l) } f'(x) = -\frac{x}{\sqrt{1-x^2}}$$

$$\text{m) } g'(w) = \frac{-2w}{\sqrt{1-(w^2+1)^2}} \quad \text{n) } f'(y) = \frac{\ln 2}{y \ln 3} 2^{\log_3 y} \quad \text{o) } f'(\theta) = -5 \csc^2 5\theta = -5 \cot^2 5\theta - 5$$

$$\text{p) } g'(x) = \frac{1}{2\sqrt{x+\sqrt{x}}}\left(\frac{1}{2\sqrt{x}}+1\right) \quad \text{q) } f'(\beta) = -2\beta \frac{\cos(\beta^2+1)}{\sin^2(\beta^2+1)} = -2\beta \csc(\beta^2+1) \cot(\beta^2+1)$$

$$\text{r) } m'(y) = \frac{1}{|y| \sqrt{9y^2-1}} \quad \text{s) } P'(t) = -\frac{\cot t}{\ln 5}$$

$$2.) \quad \text{a) } \frac{-1}{6(2a-7)^3} + C \quad \text{b) } \frac{1}{2} \ln |2a-7| + C \quad \text{c) } \frac{1}{2}e - \frac{1}{2}e^{-3} \quad \text{d) } \frac{\pi}{4} \quad \text{e) } \sin^{-1} y + C \quad \text{f) } \frac{7}{\ln 2}$$

$$\text{g) } \sec^{-1} x + C \quad \text{h) } -\csc x + C \quad \text{i) } 1 - \frac{\pi}{4}$$

$$3.) \quad \text{a) } \sqrt{x} \quad \text{b) } -\sqrt{x} \quad \text{c) } 27\sqrt{x} \quad \text{d) } \frac{1}{1+x^2} \quad \text{e) } \frac{2x}{1+x^4} \quad \text{f) } \frac{2 \tan^{-1} x}{1+x^2}$$

For more documents like this, visit our page at <https://teaching.martahidegkuti.com> and click on Lecture Notes.
E-mail questions or comments to mhidegkuti@ccc.edu.