
The Pythagorean theorem is a fundamental theorem about the connection between the three sides of a right triangle. Let us recall first a few things we will need for this topic.

Standard labeling simplifies notation. According to standard labeling, side a is opposite point A and angle α. Similarly, side b is opposite of point B and angle β, and side c is opposite point C and angle γ.



Also recall that the three angles in any triangle add up to 180°.

In any triangle, the order between sides is the same as the order between the angles opposite them. For example, the longest side is always opposite the greatest angle, and vica versa: the greatest angle is opposite the longest side. The shortest side is opposite the smallest angle, and vica versa: the smallest angle is opposite of the shortest side. If two sides are equally long, the angles opposite them are also equal.

In a right triangle, the right angle is the single greatest angle in the triangle, because the other two angles must add up to 90° so they both are smaller than 90°.

Therefore, right triangles have a single longest side and it is located opposite of the right angle.

Definition: In a right triangle, the side opposite the right angle is the longest side. We call this side the **hypotenuse** of the triangle.

We are now ready to state the Pythagorean theorem. It actually has two parts.

Theorem: (The Pythagorean theorem)

Part 1. If ABC is a right triangle with shorter sides a and b and hypotenuse c, then

$$a^2 + b^2 = c^2$$

Part 2. In any triangle with sides a, b, and c, if $a^2 + b^2 = c^2$, then the angle opposite side c measures 90°.

The first part tells us how right triangles behave. The second part tells us that *only* right triangles have this behavior.

Example 1. In each case, determine whether the three sides given are sides in a right triangle or not.

a) 5cm, 7cm, and 9cm b) 5ft, 13ft and 12ft c) 53 units, 28 units, and 45 units.

Solution: We will use the second part of the Pythagorean theorem. Let us denote the longest side by c and the other two sides by a and b. If the statement $a^2 + b^2 = c^2$, is true, the triangle is a right triangle. If not, the triangle has no right angle in it.

a) The only side here that could be the hypotenuse, is the longest one, measuring 9cm. Therefore, the two quantities we need to compare are $(5\text{cm})^2 + (7\text{cm})^2$ and $(9\text{cm})^2$.

$(5\text{cm})^2 + (7\text{cm})^2 = 25\text{cm}^2 + 49\text{cm}^2 = 74\text{cm}^2$ and $(9\text{cm})^2 = 81\text{cm}^2$

Since $74\text{cm}^2 \neq 81\text{cm}^2$, this triangle is not a right triangle.

© Hidegkuti, 2023

b) The only side here that could be the hypotenuse, is the longest one, measuring 13ft. Therefore, the two quantities we need to compare are $(5\text{ft})^2 + (12\text{ft})^2$ and $(13\text{ft})^2$.

$(5\text{ft})^2 + (12\text{ft})^2 = 25\text{ft}^2 + 144\text{ft}^2 = 169\text{ft}^2$ and $(13\text{ft})^2 = 169\text{ft}^2$

Since $169\text{ft}^2 = 169\text{ft}^2$, this triangle is a right triangle with hypotenuse 13ft long.

c) The only side here that could be the hypotenuse, is the longest one, measuring 53 units. Therefore, the two quantities we need to compare are $28^2 + 45^2$ and 53^2.

$28^2 + 45^2 = 784 + 2025 = 2809$ and $53^2 = 2809$

Since $2809 = 2809$, this triangle is a right triangle.

If we know two sides of a right triangle, the Pythagorean theorem enables us to find the third side.

Example 2. In each case, find the missing side of the right triangle.

a) the shortest two sides are 12cm and 35cm


b) the longest two sides are 15ft and 17ft

c) the shortest two sides are both 1 unit long

Solution: a) We will use the first part of the Pythagorean theorem. Let us denote the hypotenuse by x.

We state the Pythagorean theorem for this right triangle.

$$12^2 + 35^2 = x^2$$
$$144 + 1225 = x^2$$
$$1369 = x^2$$
$$\pm 37 = x$$




Since x represents a distance, and distances can never be negative, $x = -37$ is ruled out as a possible solution. Thus the hypotenuse of this triangle is 37cm.

b) We will use the first part of the Pythagorean theorem. Let us denote the missing shortest side by x.

We state the Pythagorean theorem for this right triangle.

$$x^2 + 15^2 = 17^2$$
$$x^2 + 225 = 289$$
$$x^2 = 64$$
$$x = \pm 8$$




Since distances cannot be negative, $x = -8$ is ruled out as a possible solution. Thus the missing side is 8ft long.

c) We will use the first part of the Pythagorean theorem. Let us denote the hypotenuse by x.

We state the Pythagorean theorem for this right triangle.

$$1^2 + 1^2 = x^2$$
$$1 + 1 = x^2$$
$$2 = x^2$$
$$\pm\sqrt{2} = x$$




Since x represents a distance, and distances can never be negative, $x = -\sqrt{2}$ is ruled out as a possible solution. Thus the hypotenuse of this triangle is $\boxed{\sqrt{2} \text{ unit}}$.

This last example shows that right triangles often have side lengths that are irrational numbers. This particular triangle is responsible for bringing $\sqrt{2}$ into the mathematics of the ancient Greeks. They did not have the concept of irrational numbers yet, and so couldn't really deal with $\sqrt{2}$.

In order to find missing sides, we do not always need to know the lengths of two sides if other information is given.

Example 3. The hypotenuse of a right triangle is 10 units long. The difference between the other two sides is 2 units. Find the missing sides of the right triangle.

Solution: If we label the shortest side by x, then the other missing side can be denoted by $x + 2$. We state the Pythagorean theorem for this right triangle and solve the equation for x.



$x^2 + (x+2)^2$	=	10^2	expand $(x+2)^2$
$x^2 + x^2 + 4x + 4$	=	100	combine like terms
$2x^2 + 4x + 4$	=	100	subtract 100
$2x^2 + 4x - 96$	=	0	factor out 2
$2\left(x^2 + 2x - 48\right)$	=	0	factor
$2\,(x+8)\,(x-6)$	=	0	


By the zero product rule, $x = -8$ or $x = 6$. Since distances can never be negative, -8 is ruled out as a possible solution. So $x = 6$. The other side, denoted by $x + 2$ must be $6 + 2 = 8$ units long. Thus, this right triangle has sides $\boxed{6, 8, \text{ and } 10 \text{ units}}$ long. We check: $6^2 + 8^2 = 36 + 64 = 100 = 10^2$, thus the triangle is indeed right. Also, $8 - 6 = 2$, so our answer is correct.

The following example will be a similar application problem, but the computation will be entirely different.

© Hidegkuti, 2023

Example 4. The shortest side of a right triangle is 20 units long. The difference between the other two sides is 8 units. Find the missing sides of the right triangle.

Solution: If we label the shorter missing side by x, then the other missing side, the hypotenuse can be denoted by $x + 8$. We state the Pythagorean theorem for this right triangle and solve the equation for x.




$$\begin{aligned} 20^2 + x^2 &= (x+8)^2 && \text{expand } (x+8)^2 \\ 400 + x^2 &= x^2 + 16x + 64 && \text{subtract } x^2 \\ 400 &= 16x + 64 && \text{subtract 64} \\ 336 &= 16x && \text{divide by 16} \\ 21 &= x \end{aligned}$$

Therefore, the shorter missing side, denoted by x is 21 units long, and the hypotenuse, denoted by $x + 8$ must be $21 + 8 = 29$ units long. So the three sides of this right triangle are $\boxed{\text{20, 21, and 29 units}}$ long. We check: the difference between 29 and 21 is indeed 8. For the Pythagorean theorem, $20^2 + 21^2 = 400 + 441 = 841$ and $29^2 = 841$, and so $20^2 + 21^2 = 29^2$. This means that this is indeed a right triangle, and so our answer is correct.

This problem turned out to be much easier because the equation was linear after x^2 was subtracted from both sides.

Example 5. The sides of an isosceles triangle are 4 units, 8 units, and 8 units long. Find the exact value of the length of the height drawn to the 4 units long side.

Solution: In case of isosceles triangles, the height drawn to the base splits the triangle into two identical right triangles. The height now can be easily computed via the Pythagorean theorem applied to one of these right triangles. It is important to recognize that the missing side in the right triangle is not the hypotenuse.




$$\begin{aligned} 2^2 + x^2 &= 8^2 \\ 4 + x^2 &= 64 \\ x^2 &= 60 \\ x &= \pm\sqrt{60} = \pm 2\sqrt{15} \implies x = 2\sqrt{15} \end{aligned}$$

Again, the negative solution of the equation is ruled out because distances can not be negative. The height belonging to the base is $2\sqrt{15}$ units long. Because we are asked for the exact value,we must present our answer in radical form, as $\boxed{2\sqrt{15} \text{ units}}$.

© Hidegkuti, 2023


Example 6. Consider the square-based pyramid shown on the picture. Its base has sides 10m (meters) long. The other faces of the pyramid are isosceles triangles with sides 10m, 12m, and 12m long.



a) Find the exact value of h, the length of the height in a triangular face.

b) Find the exact value of H, the length of the height of the pyramid.

Solution: a) Let us label some of the vertices as shown on the figure below. We can find the value of h by stating the Pythagorean theorem on the right triangle ABD. The hypotenuse is the side opposite the right angle.



$$\begin{aligned} (5\text{m})^2 + h^2 &= (12\text{m})^2 \\ 25\text{m}^2 + h^2 &= 144\text{m}^2 \\ h^2 &= 119\text{m}^2 \\ h &= \pm\sqrt{119\text{m}^2} \implies h = \sqrt{119}\text{m} \end{aligned}$$

Therefore, the height of the triangular face is $\boxed{\sqrt{119}\text{m}}$. This is also called **the slant height** of the pyramid.

b) Let use the labels shown on the figure above. We can find the value of H by stating the Pythagorean theorem on the right triangle BCD. The hypotenuse is the side opposite the right angle.

$$\begin{aligned} (5\text{m})^2 + H^2 &= h^2 \\ (5\text{m})^2 + H^2 &= \left(\sqrt{119}\text{m}\right)^2 \\ 25\text{m}^2 + H^2 &= 119\text{m}^2 \\ H^2 &= 94\text{m}^2 \\ H &= \pm\sqrt{94\text{m}^2} \implies H = \sqrt{94}\text{m} \end{aligned}$$


Thus the height of the pyramid is $\boxed{\sqrt{94}\text{m}}$ long.

© Hidegkuti, 2023

Example 7. A square is written into a circle of radius 5cm, i.e. all four vertices of the square are on the circle. Find the exact area of the shaded region, the part of the circle outside of the square.

Solution: The area of the shaded region is the area of the circle minus the area of the square. The area of the circle is $\pi r^2 = \pi \cdot 5^2 = 25\pi \text{cm}^2$.


To find the area of the square, we draw a diagonal of the square. The diagonal passes through the center of the circle, so it is a diameter, 10cm long. The diagonal splits the square into two right triangles. Let x denote the sides of the square. We state the Pythagorean theorem for one of these right triangles.



$$\begin{aligned} x^2 + x^2 &= 10^2 \\ 2x^2 &= 100 \\ x^2 &= 50 \end{aligned}$$


We do not even need to find x: the area of the square is x^2, so it is 50cm^2. Therefore, the area of the shaded region is $\boxed{(25\pi - 50)\ \text{cm}^2}$.

Example 8. The picture shows two identical semicircles written into a third semicircle. If the identical semicircles are of radius 1, what is the exact value of the radius of the circle?



Solution: Let us label the unknown radius by x. If two circles are tangent to each other, then the line connecting the two centers will always include the point of tangency.

Based on that, we connect the centers of circles or semicircles tangent to each other. Let us first observe that the radius of the greatest semi circle is 2 units. We will find x by solving the equation we obtain by stating the Pythagorean theorem for triangle PCD. $\overline{CD} = 1$, $\overline{DP} = x + 1$, and $\overline{PC} = 2 - x$.



Therefore,
$$\begin{aligned} 1^2 + (2-x)^2 &= (x+1)^2 \\ x^2 - 4x + 5 &= x^2 + 2x + 1 \\ -4x + 4 &= 2x \\ 4 &= 6x \\ x &= \frac{2}{3} \end{aligned}$$

So the radius of the circle is $\boxed{\frac{2}{3}\ \text{units}}$.

© Hidegkuti, 2023




Sample Problems


1. Could the three line segments given below be the three sides of a right triangle? Explain your answer.

   a) 6cm, 10cm, and 8cm b) 7ft, 15ft, and 50ft c) 4m, 5m, and 6m

2. Find the hypotenuse of the triangle shown on the figure.




3. Find the missing leg of the right triangle shown on the picture.



4. The sides of an isosceles triangle are 42 units, 29 units, and 29 units long. Find the length of the height drawn to the 42 units long side.

5. The hypotenuse of a right triangle is 20cm. The difference between the other two sides is 4cm. Find the sides of the triangle.




6. Find the length of the longest line segment inside the rectangular prism shown on the picture. (This line segment is called the **main diagonal** of the prism.)

7. Find the height h of the cone shown on the picture, if the base has a radius of $r = 10$m and $a = 26$m.

8. An arch is in the shape of a semicircle. At a point along the base 1 foot from an end of the arch, the height of the arch is 7 feet. Find the maximum height of the arch.

9. C is the center of a circle. P is a point outside of the circle. We drew a tangent line to the circle from point P. The point of tangency is Q. The distance between P and C is 34 units, and the distance between P and Q is 30 units. How long is the radius of the circle?




© Hidegkuti, 2023


10. A pyramid (shown on the picture) has a square base with sides 10m (meters) long. The other faces of the pyramid are isosceles triangles with sides 10m, 12m, and 12m.

    a) Find the exact value of h, the length of the height in a triangular face.

    b) Find the exact value of H, the length of the height of the pyramid.



11. Four identical circles are arranged in a fifth circle as shown on the picture. Compute the exact value of the radius of the big circle if the smaller circles have radius 1.






Practice Problems


1. Could the three line segments given below be the three sides of a right triangle? Explain your answer.

   a) 3 cm, 7 cm, and 8cm b) 37ft, 12ft, and 35ft c) 6m, 7m, and 8m

2. Find the hypotenuse of the triangle shown on the figure.



3. Find the missing leg of the right triangle shown on the picture.



4. Find the length of the diagonal in a rectangle with sides 20 ft and 21 ft long.

5. Find the length of the diagonal of a square with sides 1 unit long.

6. The sides of an isosceles triangle are 25m, 25m, and 14m long. Find the length of the height drawn to the 14m long side.

7. Two sides of a right triangle are 8cm and 17cm long. Find the length of the missing side.

8. One leg of a right triangle is 9cm. The difference between the other two sides is 1cm. Find the length of all sides.

© Hidegkuti, 2023

9. The hypotenuse of a right triangle is 25 in. The difference between the other two sides is 17 in. Find the length of all sides.


10. The shortest side of a right triangle is 16 units long. The difference between the other two sides is 4 units. Find the sides of this right riangle.

11. Consider a triangle with sides 7 unit, 7 unit, and 12 units long.


    a) Compute the exact value of the height drawn to the longest side.

    b) Compute the exact value of the area of the triangle.


12. Find the missing lengths indicated on the picture below.



a)



b)



c)

13. Find the length of the main diagonal in a rectangular prism with sides

    a) 2m, 10m, and 11m b) 5ft, 7ft, and 1ft

14. The height of a regular (same as equilateral) triangle is 6 feet. How long are its sides? Give the exact value of the answer.

15. The diagonal of a square is 80 meters long. How long are the sides?

16. Suppose that C is a center of a circle and P is a point (outside of the circle) 9 units away from C. We draw a tangent line to the circle from point P. Let Q be the point of tangency. Given that line segment PQ is 8 units long, compute the exact value of

    a) the radius of the circle. b) the area of triangle CPQ.


17. An arch is in the shape of a semicircle. At a point along the base 2 meters from an end of the arch, the height of the arch is 10 meters. Find the maximum height of the arch.

18. How long is the main diagonal in a cube of sides 1 meter long?


© Hidegkuti, 2023

19. Find the exact value of the missing lengths, labeled

a) x and y




b) p and q




20. Consider a pyramid with a square base. The base has sides each 20 meters long. All other edges are 20 meters long. Compute the exact value of the height of the pyramid.


21. a) Compute the area of a square written into a semicircle with radius 1.



b) Compute the area of a square written into a circle with radius 1.




22. We drew a semicircle into an equilateral triangle as shown in the picture. (The center of the circle is the midpoint of one side, and the other two sides are tangents to the circle.) Find the radius if the triangle has sides 10 units long.



23. Consider a regular triangle width sides 1 unit long, written into a circle as shown on the picture.

a) Find the exact value of the radius of the circle.


b) The center of the circle splits the height of the traingle into two parts. What is the ratio of the shorter part to the longer part?




24. Square $ABCD$ is written in a circle of radius 10 units as follows. Points A and B are on the circle, and side CD is tangent to the circle. How long are the sides of the square?

© Hidegkuti, 2023


25. A square is written into a circle of radius 3 inches. Find the exact area of the shaded region, the part of the circle outside of the square.



26. (Enrichment) Four squares are shown on the picture. The radius of the quarter circle is 10 units. Find the area of the shaded (blue) region. (problem from Catriona Agg)



27. (Enrichment) Two spheres are resting against the wall as shown on the picture. The radii of the spheres are 5 feet and 2 feet long. How high is the center of the smaller sphere?



© Hidegkuti, 2023



Answers

Sample Problems


1. a) There is a right angle opposite the 10cm long side. b) not a right triangle 2. 17m 3. 35in
4. 20 units 5. 12cm and 16cm 6. $\sqrt{45}$ft 7. 24m 8. 25ft 9. 16 units
10. $h = \sqrt{119}$m $H = \sqrt{94}$m 11. $1 + \sqrt{2}$

Practice Problems

1. a) not a right triangle b) There is a right angle opposite the 37ft long side. c) not a right triangle
2. 13mi 3. 16cm 4. 29ft 5. $\sqrt{2}$ unit 6. 24m 7. 15cm or $\sqrt{353}$cm
8. 9cm, 40cm, and 41cm 9. 7in, 24in, and 25in
10. 16, 30, and 34 units 11. a) $\sqrt{13}$ unit b) $6\sqrt{13}$ unit^2 12. a) 20 b) 7 c) 10
13. a) 15m b) $\sqrt{75}$ft 14. $4\sqrt{3}$ft 15. $40\sqrt{2}$m 16. a) $\sqrt{17}$ unit b) $4\sqrt{17}$ unit^2
17. 26m 18. $\sqrt{3}$m 19. a) $x = \sqrt{105}$ft $y = \sqrt{89}$ft b) $q = 25$cm, $p = \sqrt{850}$cm $= 5\sqrt{34}$cm
20. $10\sqrt{2}$m 21. a) $\frac{4}{5}$ unit^2 b) 2 unit^2 22. $\frac{5\sqrt{5}}{2}$ units
23. a) $R = \frac{1}{\sqrt{3}} = \frac{\sqrt{3}}{3}$ b) 1 to 2 24. 16 units 25. $(9\pi - 18)$ in^2 26. 68 unit^2 27. $2\sqrt{10}$ feet

© Hidegkuti, 2023

Sample Problems



1. Could the three line segments given be the three sides of a right triangle? Explain your answer.

   a) 6 cm, 10 cm, and 8 cm

   Solution: The longest side is 10 cm long. Thus, only this side can be the hypotenuse. We use the Pythagorean theorem to check for a right angle:

   $$6^2 + 8^2 \stackrel{?}{=} 10^2$$

   We get that the two quantities are equal, thus this triangle has a right angle.

   b) 4 m, 5 m, and 6 m

   Solution: The longest side is 6 m long. Thus, only this side can be the hypotenuse. We use the Pythagorean theorem to check for a right angle: (LHS is short for left-hand side and RHS for right-hand side.)

   $$4^2 + 5^2 \stackrel{?}{=} 6^2$$


   $$\text{LHS} = 4^2 + 5^2 = 16 + 25 = 41 \qquad \text{RHS} = 6^2 = 36 \qquad \text{LHS} \neq \text{RHS}$$

   We get that the two quantities are not equal, thus this triangle does not have a right angle.

2. Find the hypotenuse of the triangle shown on the figure.

   Solution: We apply the Pythagorean theorem. The longest side is always the one opposite the right angle.

   $$\begin{aligned} 8^2 + 15^2 &= x^2 \\ 289 &= x^2 \\ \pm 17 &= x \end{aligned}$$

   Since distance can not be negative, -17 is ruled out. The answer is 17m.


   Note that the equation $x^2 = 289$ has two solutions, 17 and -17. In the context of the geometry, only the positive solution is possible.

© Hidegkuti, 2023

3. Find the missing leg of the right triangle shown on the picture.

   Solution: We apply the Pythagorean theorem. The longest side is always the one opposite the right angle.

$$
\begin{aligned}
(12\text{in})^2 + x^2 &= (37\text{in})^2 \\
x^2 + 144\text{in}^2 &= 1369\text{in}^2 \qquad \text{subtract } 144\text{in}^2 \\
x^2 &= 1225\text{in}^2 \qquad \sqrt{1225} = 35 \\
x &= \pm 35\text{in}
\end{aligned}
$$




   Since distance can not be negative, -35in is ruled out. The answer is 35 inches.

4. The sides of an isosceles triangle are 42 units, 29 units, and 29 units long. Find the length of the height drawn to the 42 units long side.

   Solution: In case of isosceles triangles, the height drawn to the base splits the triangle into two identical right triangles as shown on the picture. The height now can be easily computed via the Pythagorean theorem.


$$\begin{aligned} 21^2 + h^2 &= 29^2 \\ 441 + h^2 &= 841 \\ h^2 &= 400 \\ h &= \pm 20 \implies h = 20 \end{aligned}$$



   Again, the negative solution of the equation is ruled out because distances cannot be negative. The height belonging to the base is 20 units long.

5. The hypotenuse of a right triangle is 20cm. The difference between the other two sides is 4cm. Find the sides of the triangle.

   Solution: Let x denote the shortest side. Then the other missing side is $x + 4$cm long. We state the Pythagorean theorem for the triangle and solve the quadratic equation for x.




$$\begin{aligned} x^2 + (x+4)^2 &= 20^2 && \text{expand } (x+4)^2 \\ x^2 + x^2 + 8x + 16 &= 400 && \text{combine like terms} \\ 2x^2 + 8x + 16 &= 400 && \text{subtract 400} \\ 2x^2 + 8x - 384 &= 0 && \text{factor out 2} \\ 2\left(x^2 + 4x - 192\right) &= 0 && \text{factor} \\ 2\,(x+16)\,(x-12) &= 0 \end{aligned}$$

   By the zero product rule, $x = -16$ or $x = 12$. Since distances are never negative, -16 is ruled out. If the shortest side is 12cm, the other side is
   12cm + 4cm = 16cm. Thus the solution is 12cm and 16cm. We check:

$$16 - 12 = 4 \checkmark \text{ and } 16^2 + 12^2 = 256 + 144 = 400 = 20^2 \checkmark$$

© Hidegkuti, 2023

6. Find the length of the longest line segment inside the rectangular prism shown on the picture. (This line segment is called the **main diagonal** of the prism.)




Solution: We will apply the Pythagorean theorem twice.

Let us label the points and sides we will use on the picture first.

We will find x using the Pythagorean theorem in triangle ABC.

Then we can find y using the Pythagorean theorem in triangle ACD.



$$\begin{aligned} 4^2 + 5^2 &= x^2 \\ 41 &= x^2 \\ x &= \pm\sqrt{41} \quad \Longrightarrow x = \sqrt{41} \end{aligned}$$


$$\begin{aligned} 2^2 + x^2 &= y^2 \\ 45 &= y^2 \\ y &= \pm\sqrt{45} \quad \Longrightarrow y = \sqrt{45} \end{aligned}$$

Note: Our result is actually $\sqrt{2^2 + 4^2 + 5^2}$. Indeed, we can see that the length of the main diagonal in a rectangular prism.with sides x, y, and z is $L = \sqrt{x^2 + y^2 + z^2}$. This is sometimes called the 3-dimensional Pythagorean theorem.

7. Find the height h of the cone shown on the picture, if the base has a radius of 10m and $a = 26$m.


Solution: There is a right triangle formed by a, h, and r as the picture shows.We state the Pythagorean theorem for this triangle and solve for h.

$$\begin{aligned} r^2 + h^2 &= a^2 \\ 10^2 + h^2 &= 26^2 \\ h^2 + 100 &= 676 \\ h^2 &= 576 \\ h &= \pm 24 \end{aligned}$$



The negative value is ruled out, and so the height is 24m.

8. An arch is in the shape of a semicircle. At a point along the base 1 foot from an end of the arch, the height of the arch is 7 feet. Find the maximum height of the arch.




Solution: Consider the picture shown. Let r denote the radius of the semi-circle. We are asked to find the value of r. $AD = 1$ and $AB = 7$ were given.
Clearly $BC = r$, and this problem becomes easy once we realize that $AC = r - 1$ since the entire line segment CD is the radius r. Now we state the Pythagorean theorem on the right triangle ABC and solve for r.

$$\begin{aligned} (r-1)^2 + 7^2 &= r^2 \\ r^2 - 2r + 1 + 49 &= r^2 \quad \text{subtract } r^2 \\ -2r + 50 &= 0 \quad \text{add } 2r \\ 50 &= 2r \quad \text{divide by 2} \\ 25 &= r \end{aligned}$$

Thus the maximum height of the arch is 25 feet.

9. C is the center of a circle. P is a point outside of the circle. We drew a tangent line to the circle from point P. The point of tangency is Q. The distance between P and C is 34 units, and the distance between P and Q is 30 units. How long is the radius of the circle?



Solution: Recall that in case of circles, the tangent line is perpendicular to the radius drawn to the point of tangency. We state the Pythagorean theorem for triangle PQC and solve for r.

$$\begin{aligned} r^2 + 30^2 &= 34^2 \\ r^2 &= 34^2 - 30^2 \\ r^2 &= 256 \\ r &= \pm 16 \end{aligned}$$


Distances cannot be negative, so we rule out -16. So the radius of the circle is 16 units long.

© Hidegkuti, 2023

10. A pyramid (shown on the picture) has a square base with sides 10m (meters) long. The other faces of the pyramid are isosceles triangles with sides 10m, 12m, and 12m.

    a) Find the exact value of h, the length of the height in a triangular face.

    b) Find the exact value of H, the length of the height of the pyramid.



Solution: Let us label some of the vertices as shown on the figure.
We can find the value of h by stating the Pythagorean theorem on the right triangle ABD. The hypotenuse is the side opposite the right angle.


$$\begin{aligned} (5\text{m})^2 + h^2 &= (12\text{m})^2 \\ 25\text{m}^2 + h^2 &= 144\text{m}^2 \\ h^2 &= 119\text{m}^2 \\ h &= \pm\sqrt{119\text{m}^2} \implies h = \sqrt{119}\text{m} \end{aligned}$$

b) Find the exact value of H, the length of the height of the pyramid.


Solution: Let use the labels shown on the figure above. We can find the value of H by stating the Pythagorean theorem on the right triangle BCD. The hypotenuse is the side opposite the right angle.

$$\begin{aligned} (5\text{m})^2 + H^2 &= h^2 \\ (5\text{m})^2 + H^2 &= \left(\sqrt{119}\text{m}\right)^2 \\ 25\text{m}^2 + H^2 &= 119\text{m}^2 \\ H^2 &= 94\text{m}^2 \\ H &= \pm\sqrt{94\text{m}^2} \quad \implies \quad H = \sqrt{94}\text{m} \end{aligned}$$

11. Four identical circles are arranged in a fifth circle as shown on the picture. Compute the exact value of the radius of the big circle if the smaller circles have radius 1.



Solution: Let us first draw a picture with the centers of all five circles indicated as shown on the picture. Please note that PQ means the line segment determined by points P and Q; and $\overline{PQ}$ means the length of that line segment.
Let R denote the radius of the larger circle and A, B, C, and D denote the centers of the small circles as shown. If we extend line segment AC to intersect the big circle, we obtain the points P and Q: Line segment PQ is the diagonal of the larger circle, and so $\overline{PQ} = 2R$. Line segment PQ is also the sum of three line segments:



$$PQ = PA + AC + CQ \text{ and so } \overline{PQ} = 2R = \overline{PA} + \overline{AC} + \overline{CQ}$$

Line segments PA and CQ are 1 unit long because they are each a radius in a smaller circle. We will compute $\overline{AC}$ by the Pythagorean theorem, stated for triangle ABC. The length of ine segments AB and BC are both 2 units long because they measure the radius twice. So

$$\begin{aligned} 2^2 + 2^2 &= \left(\overline{AC}\right)^2 \\ 8 &= \left(\overline{AC}\right)^2 \\ \pm\sqrt{8} &= \overline{AC} \qquad \Longrightarrow \quad \overline{AC} = \sqrt{8} = 2\sqrt{2} \end{aligned}$$

In this problem it is important that $\sqrt{8}$ can be simplified using the identity $\sqrt{ab} = \sqrt{a}\sqrt{b}$.

$$\sqrt{8} = \sqrt{4 \cdot 2} = \sqrt{4} \cdot \sqrt{2} = 2\sqrt{2}$$

We are now ready to compute R.

$$\begin{aligned} 2R &= \overline{PA} + \overline{AC} + \overline{CQ} \\ 2R &= 1 + 2\sqrt{2} + 1 \\ 2R &= 2 + 2\sqrt{2} \\ R &= \frac{2 + 2\sqrt{2}}{2} = \frac{2\left(1 + \sqrt{2}\right)}{2} = 1 + \sqrt{2} \end{aligned}$$

And so $\boxed{R = 1 + \sqrt{2}}$.

For more documents like this, visit our page at https://teaching.martahidegkuti.com and click on Lecture Notes. E-mail questions or comments to mhidegkuti@ccc.edu.

© Hidegkuti, 2023